

REPORT DOCUMENTATION PAGEForm Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.

1. REPORT DATE (DD-MM-YYYY) 22-05-2001		2. REPORT TYPE Journal Article		3. DATES COVERED (From - To) Winter 2000	
4. TITLE AND SUBTITLE Nowcasting and Forecasting The Global Ocean				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER PE623207N/ PE62435N	
				5d. PROJECT NUMBER	
6. AUTHOR(S) E. Joseph Metzger, Ole Martin Smedstad, Harley E. Hurlburt, Alan J. Wallcraft and Robert C. Rhodes				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Research Laboratory Oceanography Division Stennis Space Center, MS 39529-5004				8. PERFORMING ORGANIZATION REPORT NUMBER JA/7320-00-1026	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) Office of Naval Research 800 N. Quincy St. Arlington, VA 22217-5660				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release, distribution is unlimited					
13. SUPPLEMENTARY NOTES Backscatter; Observing Aquatic Environments Winter 2000					
14. ABSTRACT Over the past decade, the Naval Research Laboratory (NRL) has been actively working on the problem of eddy-resolving global ocean modeling and prediction. The world's first global ocean nowcast/forecast system has been developed at NRL and is running in real-time at the Naval Oceanographic Office (NAVO). The system consist of the 1/16 degree 7-layer, thermodynamic, finite depth version of the Navy Layered Ocean Model (NLOM) for the global ocean (72 degrees to 65 degrees north) and includes a mixed layer and sea surface temperature (SST). It spun-up to real-time using high frequency wind and thermal forcing from the Fleet Numerical Meteorology and Oceanography Center's Navy Operational Global Atmospheric Prediction system. It includes assimilation of SST and near real-time TOPEX/POSEIDON and ERS-2 altimeter data made available from NAVO's Altimeter Data Fusion Center NAVO is scheduled to begin operational testing of the system on 15 December 2000 and pending a successful outcome it will become an operational product. system gives a real-time view of the ocean down to 50-200 km scale of ocean eddies and the meandering of ocean currents and fronts. Figure 1 is an example of sea surface height (SSH) and SST for the Gulf Stream in the Atlantic.					
15. SUBJECT TERMS nowcasting and forecasting, eddy-resolving, meandering					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT Unclassified	b. ABSTRACT Unclassified	c. THIS PAGE Unclassified	SAR	4	E. Joseph Meztger
					19b. TELEPHONE NUMBER (include area code) (228) 688-4762

20010808 035

PUBLICATION OR PRESENTATION RELEASE REQUEST

SSC-200-00

Division 7300 Pub Key: 2692

1. REFERENCES AND ENCLOSURES Ref: (a) NRL Instruction 5600.2 (b) NRL Instruction 5510.40D Encl: (1) Two copies of subject paper (or abstract)	2. TYPE OF PUBLICATION OR PRESENTATION <input type="checkbox"/> Abstract only, published <input type="checkbox"/> Book <input type="checkbox"/> Conference Proceedings (refereed) <input type="checkbox"/> Invited speaker <input type="checkbox"/> Journal article (refereed) <input type="checkbox"/> Oral Presentation, published <input checked="" type="checkbox"/> Other explain <u>Quarterly Magazine</u> <input type="checkbox"/> Abstract only, not published <input type="checkbox"/> Book Chapter <input checked="" type="checkbox"/> Conference Proceedings (not refereed) <input type="checkbox"/> Multimedia report <input type="checkbox"/> Journal article (not refereed) <input type="checkbox"/> Oral Presentation, not published	3. ADMINISTRATIVE INFORMATION STRN <u>547320--00-1026</u> Route Sheet No. _____ Job Order No. <u>73-5094-01</u> Classification <u>X</u> <u>u</u> <u>C</u> Sponsor <u>ONR/SPAWAR</u> approval obtained <u>X</u> yes _____
---	---	---

4. AUTHOR	
Title of Paper or Presentation <u>Nowcasting & Forecasting the Global Ocean</u> <u>An Operational 1/16 degree Global Ocean Nowcast/Forecast System (changed title)</u>	
Author(s) Name(s) (First, Mi, Last), Code, Affiliation if not NRL <u>E.J. Metzger, Code 7323; O.M. Smedstad, PSI; H.E. Hurlburt, A.J. Wallcraft, R.C. Rhodes, Code 7323</u>	
It is intended to offer this paper to the <u>Backscatter - Official Magazine of the Alliance for Marine Remote Sensing Association</u> (Name of Conference) <u>Winter 2000</u> (Date, Place and Classification of Conference)	
and/or for publication in _____ (Name and Classification of Publication) (Name of Publisher)	
After presentation or publication, pertinent publication/presentation data will be entered in the publications data base, in accordance with reference (a). It is the opinion of the author that the subject paper (is _____) (is not <u>X</u>) classified, in accordance with reference (b). This paper does not violate any disclosure of trade secrets or suggestions of outside individuals or concerns which have been communicated to the Laboratory in confidence. This paper (does _____) (does not <u>X</u>) contain and militarily critical technology. This subject paper (has _____) (has never <u>X</u>) been incorporated in an official NRL Report.	
<u>E. Joseph Metzger, Code 7323</u> Name and Code (Principal Author) <u>E. Joseph Metzger</u> (Signature)	

CODE	SIGNATURE	DATE	COMMENTS
Author(s) E.J. Metzger, Code 7323	<u>E. Joseph Metzger</u>	<u>12/13/2000</u>	
Section Head Rhodes or Acting	<u>Robert Rhodes</u>	<u>12/13/00</u>	
Branch Head Harding or Acting	<u>John Harding</u>	<u>12/13/00</u>	
Division Head	<u>David B. Corman</u>	<u>1/16/01</u>	1. Release of this paper is approved. 2. To the best knowledge of this Division, the subject matter of this paper (has _____) (has never <u>X</u>) been classified.
Payne or Acting	<u>David B. Corman</u>	<u>1/16/01</u>	1. Paper or abstract was released. 2. A copy is filed in this office. <u>SSC-200-00</u>
Security, Code 7030.1	<u>David B. Corman</u>	<u>1/16/2001</u>	
Office of Counsel, Code 1008.3	<u>David B. Corman</u>	<u>1/16/2001</u>	
ADOR/Director NCST	<u>David B. Corman</u>	<u>1/16/2001</u>	
Public Affairs (Unclassified/ Unlimited Only), Code 7030.4	<u>David B. Corman</u>	<u>1/16/2001</u>	
Division, Code			
Author, Code			

Hex 2692

6. DISTRIBUTION STATEMENTS (Author to check appropriate statement and fill reason as required)

☒ **A - Approved for public release, distribution is unlimited.**

☐ **B - Distribution authorized to U.S. Government agencies only (check reason below):**

- | | | |
|---|--|--|
| <input type="checkbox"/> Foreign Government Information | <input type="checkbox"/> Contractor Performance Evaluation | <input type="checkbox"/> Critical Technology |
| <input type="checkbox"/> Proprietary Information | <input type="checkbox"/> Administrative/Operational Use | <input type="checkbox"/> Premature Dissemination |
| <input type="checkbox"/> Test and Evaluation | <input type="checkbox"/> Software Documentation | <input type="checkbox"/> Cite "Specific Authority" _____ |

Date statement applied _____

(Identification of valid documented authority)

Other requests for this document shall be referred to _____

(Insert Controlling DOD Office*)

☐ **C - Distribution authorized to U.S. Government agencies and their contractors (check reason below):**

- | | | |
|---|---|--|
| <input type="checkbox"/> Foreign Government Information | <input type="checkbox"/> Software Documentation | <input type="checkbox"/> Cite "Specific Authority" _____ |
| <input type="checkbox"/> Administrative/Operational Use | <input type="checkbox"/> Critical Technology | |

Date statement applied _____

(Identification of valid documented authority)

Other requests for this document shall be referred to _____

(Insert Controlling DOD Office*)

☐ **D - Distribution authorized to DOD and DOD contractors only (check reason below):**

- | | | |
|---|--|--|
| <input type="checkbox"/> Foreign Government Information | <input type="checkbox"/> Critical Technology | <input type="checkbox"/> Cite "Specific Authority" _____ |
| <input type="checkbox"/> Software Documentation | | |
| <input type="checkbox"/> Administrative/Operational Use | | |

Date statement applied _____

(Identification of valid documented authority)

Other requests for this document shall be referred to _____

(Insert Controlling DOD Office*)

☐ **E - Distribution authorized to DOD components only (check reason below):**

- | | | |
|---|--|--|
| <input type="checkbox"/> Proprietary Information | <input type="checkbox"/> Premature Dissemination | <input type="checkbox"/> Critical Technology |
| <input type="checkbox"/> Foreign Government Information | <input type="checkbox"/> Software Documentation | <input type="checkbox"/> Direct Military Support |
| <input type="checkbox"/> Administrative/Operational Use | <input type="checkbox"/> Contractor Performance Evaluation | <input type="checkbox"/> Test and Evaluation |
| | | <input type="checkbox"/> Cite "Specific Authority" _____ |

Date statement applied _____

(Identification of valid documented authority)

Other requests for this document shall be referred to _____

(Insert Controlling DOD Office*)

☐ **F - Further dissemination only as directed by _____**

(Insert Controlling DOD Office*)

Date statement applied _____ or higher DOD authority _____

☐ **X - Distribution authorized to U.S. Government agencies and private individuals or enterprises eligible to obtain export-controlled technical data in accordance with regulations implementing 10 U.S.C. 140c.**

Date statement applied _____


Other requests for this document shall be referred to _____

(Insert Controlling DOD Office*)

*For NRL publications, this is usually the Commanding Officer, Naval Research Laboratory, Washington, DC 20375-5320

7. OTHER LIMITATION

- ☐ Classification ☐ NOFORN ☐ DTIC exempt (explain) _____


Classification Review
(initial/Date)

Substantive changes made in this document after approval by Classification Review and Public Release invalidate these reviews. Therefore, if any substantive changes are made by the author, Technical Information, or anyone else, the document must be returned for another Classification Review and Public Release.

8. INSTRUCTIONS

Author completes and submits this form with the manuscript via line channels to the division head for review and approval according to the routing in section 4.

1. NRL Reports Submit the diskette (if available), manuscript, typed double-spaced, complete with tables, illustrations, references, draft SF 298, and proposed distribution list.
2. NRL Memorandum Reports Submit a copy of the original, typed manuscript complete with tables, illustrations, references, draft SF 298, and proposed distribution list.
3. NRL Publications or other books, brochures, pamphlets, Handled on a per case basis by Site Technical Information Office. proceedings, or any other printed publications.



Nowcasting & Forecasting the Global Ocean

E. Joseph Metzger (1), Ole Martin Smedstad (2), Harley E. Rurlburt (1), Alan J. Wallcraft (1), Robert C. Rhodes (1)

(1) Naval Research Laboratory, Stennis Space Center, MS, USA

(2) Planning Systems Incorporated, Stennis Space Center, MS, USA

E-mail: metzger@nrlssc.navy.mil

Over the past decade, the Naval Research Laboratory (NRL) has been actively working on the problem of eddy-resolving global ocean modeling and prediction. The world's first global ocean nowcast/forecast system has been developed at NRL and is running in real time at the Naval Oceanographic Office (NAVO).

The system consists of the 1/16° seven layer, thermodynamic, finite depth version of the Navy Layered Ocean Model (NLOM) for the global ocean (72°S to 65°N) and includes a mixed layer and sea surface temperature (SST). It was spun-up to real time using high frequency wind and thermal forcing from the Fleet Numerical Meteorology and Oceanography Center's Navy Operational Global

represents a zone of fresher water with low scattering and low chlorophyll growth. Ternary diagrams may provide a new way to classify the complex, coastal Case II waters into biogeochemical provinces.

Summary

With the availability of new ocean color sensors, the application of remote sensing for understanding coastal processes is beginning. The rich spectral signature of coastal waters allows an opportunity to separate the in-water components into tracers of processes. Improved atmospheric and in-water algorithms in bio-optical sensing have

been extended into coastal waters. Unlike open ocean waters, coastal waters have a complex variety of combinations of particles, (organic and inorganic), and absorption components (chlorophyll and colored dissolved organic matter).

Because coastal waters are more complex than open ocean waters, improved methods of classification are needed to account for a variety of processes. By basing a classification on remote sensing bio-optical products, we enhance our ability to understand the spatial variability of the processes.

References

Arnone, R. A. and R.W. Gould, Coastal Monitoring Using Ocean Color, *Sea Technology*, 39(9), 18-27, September, 1998.

Arnone, R. A., Integrating Satellite Ocean Color into Navy Operations, *Backscatter*, 10(3), 8-12, August, 1999.

Morel A. and L. Prieur, Analyses of Variations in Ocean Color. *Limnol Oceanogr*, 22, 709-722, 1977.

Sathyendrenath, S. (ed.), Remote Sensing of Ocean Colour in Coastal and Other Optically-Complex Waters. *Reports of the International Ocean Colour Coordinating Group*, No3, IOCCG, Dartmouth, Canada, 2000.

Water mass – Ternary Diagram




Figure 4. Coastal waters are complex mixtures of backscattering (550 nm), chlorophyll concentration and absorption from colored dissolved organic matter. These three SeaWiFS products are classified using a ternary diagram, which bases a water mass on the percentage of each component. The colors represent mean water mass from the scatter plot (Figure 3) and illustrate the transition of coastal to open ocean waters based on these components.




Figure 1. Snapshots of sea surface height (top) and sea surface temperature (bottom) zoomed in on the Gulf Stream region for 6 December 2000 from the real time 1/16° global nowcast/forecast system with assimilation of SST and satellite altimeter data from Topex/Poseidon and ERS-2. Superimposed on the sea surface height (SSH) is the frontal analysis based on infrared satellite imagery from the Warfighting Support Center at NAVO. Superimposed on the SST is the Gulf Stream IR northwall 1982-96 mean \pm standard deviation determined by P. Cornillon (University of Rhode Island) and Z. Sirkes (University of Southern Mississippi).

Atmospheric Prediction System. It includes assimilation of SST and near-real time Topex/Poseidon and ERS-2 altimeter data made available from NAVO's altimeter data

fusion center. NAVO began operational testing of the system last December, and pending a successful outcome, it will become an operational product. The system gives a real

time view of the ocean down to the 50-200 km scale of ocean eddies and the meandering of ocean currents and fronts. Figure 1 is an example of sea surface height (SSH) and




Figure 2. Snapshots of sea surface height (SSH) for the North Atlantic Ocean from four simulations using the Navy Layered Ocean Model at a) $1/8^\circ$, b) $1/16^\circ$, c) $1/32^\circ$, and d) $1/64^\circ$ resolution. The model is forced by the monthly wind stress climatology from Hellerman and Rosenstein (1983) and a port driven thermohaline circulation. There is no SSH assimilation in these model results. Adapted from Hurlburt and Hogan (2000).

SST for the Gulf Stream in the Atlantic.

Atmospheric Versus Oceanic Prediction

The first operational weather prediction occurred in May 1955 as a joint air force, navy and weather bureau project. In principle, numerical ocean modeling is similar to atmospheric modeling but global operational oceanography has lagged far behind because of two major complications. (1) Oceanic space and time scales are much different than those of the atmosphere. Ocean eddies are typically about 100 km in diameter which makes them 20 to 30 times smaller than comparable atmospheric highs and lows. This means that approximately four orders of magnitude more computer time and three orders of magnitude more computer memory are required. (2) Unlike the meteorological radiosonde network that provides initial conditions from the surface to near the top of the atmosphere, there are very few observations below the ocean surface at the synoptic time scale. Thus, any effective oceanic data assimilative technique will be limited to surface satellite observations. One advantage ocean modeling enjoys is that sea surface height forecast skill is longer than the 10 to 14 day limit for atmospheric pressure systems as will be shown later.

Model Resolution Requirements

A major component of NRL's ocean modeling program has been a detailed study of the resolution required for ocean prediction. There is strong evidence that the Navy Layered Ocean Model and other popular ocean models need to use grid cells for each prognostic variable that are at most about 8 km across at mid-latitudes. NRL research has shown that doubling the horizontal resolution to 4 km per cell gives substantial improvement but doubling again to 2 km gives only modest additional improvement (Hurlburt and Hogan, 2000). For the Navy Layered Ocean Model grid these resolutions translate to $1/16^\circ$, $1/32^\circ$ and $1/64^\circ$, respectively. This is for the global and basin-scale.

At 4 km, the optimal resolution is finer than might be expected based on the size of eddies. In relation to ocean eddy size it is similar to the resolution currently used by the leading weather forecasting models in relation to the size of atmospheric highs and lows. More specifically, our research has shown that fine resolution of the ocean eddy scale is required to obtain coupling between upper ocean currents and seafloor topography via turbulent flow instabilities. This coupling can strongly affect the pathways of upper ocean currents and fronts, including the Gulf Stream in the Atlantic and the Kuroshio in the Pacific. The high resolution is

also required to obtain sharp fronts that span major ocean basins. It can even affect the large scale shape of ocean gyres such as the Sargasso Sea in the Atlantic.

The need for high horizontal resolution is highlighted in Figure 2 that shows sea surface height snapshots for the Atlantic basin from four non-data assimilative Navy Layered Ocean Model simulations at $1/8^\circ$, $1/16^\circ$, $1/32^\circ$ and $1/64^\circ$ resolution. The $1/8^\circ$ model shows two unrealistic Gulf Stream pathways that are consistent with linear dynamics. Higher horizontal resolution is required in order to get into a more non-linear flow regime, and this occurs at $1/16^\circ$

resolution. Note the explosion of eddies between the $1/8^\circ$ and $1/16^\circ$ models. There is now one Gulf Stream pathway that separates from the U.S. coast at Cape Hatteras, although it does not exhibit a strong inertial character as far into the basin as is observed. In the $1/32^\circ$ model, it penetrates farther into the interior, changing the large scale shape of the subtropical gyre (Sargasso Sea) in comparison to the $1/8^\circ$ model. The $1/64^\circ$ model provides modest improvement over the $1/32^\circ$ model.

Computational Requirements

As far back as 1989, the President's Office of Science and Technology recognized




Figure 3. Zoom on the Kuroshio south and east of Japan. (a) SSH (sea surface height) for 15 January 1999 from the $1/16^\circ$ global nowcast/forecast system with assimilation of satellite altimeter data from Topex/Poseidon and ERS-2. The altimeter tracks with data available for this update cycle are overlaid. (b) The corresponding SSH snapshot from a 14 day forecast initialized from 1 January 1999. (c) The MODAS $1/8^\circ$ SST analysis from satellite IR imagery. The MODAS SST product was developed by C. Barron at NRL and is an operational product at NAVO. The SST color bar is designed to highlight the Kuroshio pathway.

amrs Corporate Members

A.U.G. Signals Ltd
www.augsignals.com

AMEC
www.amec.com

ARGOSS
www.argoss.nl

ASL Environmental Sciences Inc.
www.aslenv.com

Atlantis Scientific Inc.
www.atlsci.com

Atmospheric & Environmental Research, Inc.
www.aer.com

Canada Centre for Remote Sensing
www.ccrs.nrcan.gc.ca

Canadian Centre for Marine Communications
www.ccmc.nf.ca

Canadian Hydrographic Service
www.chs-shc.dfo-mpo.gc.ca

Canadian Ice Service
www.ec.gc.ca

Canadian Space Agency
www.space.gc.ca

Center for Coastal Physical Oceanography
www.ccpo.edu

Chelsea Instruments Limited
www.chelsea.co.uk

CODAR Ocean Sensors, Ltd.
www.codaros.com

Comserve Australia Pty Ltd
www.themaps.com.au

global ocean modeling and prediction as a "Grand Challenge" problem, defined as requiring a computer system capable of sustaining at least one trillion floating point adds or multiplies per second. We are solving the problem on today's systems capable of only a few percent of this performance by taking a multi-faceted approach to cost minimization.

One facet is the use of the Navy Layered Ocean Model (Wallcraft and Moore, 1997) which has been specifically designed for eddy-resolving global ocean prediction. It is tens of times faster than other ocean models in computer time per model year for a given horizontal resolution and model domain. The Navy Layered Ocean Model's performance is in turn due to a range of design decisions, the most important of which is the use of isopycnal (density tracking) layers in the vertical rather than the

more usual fixed depth cells. Density is the natural vertical coordinate system for the stratified ocean, and it allows seven Navy Layered Ocean Model layers to replace the 100 or more fixed levels that would be needed at $1/16^\circ$ resolution.

Another facet of our efficiency drive is the use of an inexpensive data assimilation scheme backed by a statistical technique for relating surface satellite data to subsurface fields. The statistics are from an atmospherically forced 20 year inter-annual simulation of the same ocean model, an application that requires a model with high simulation skill.

The initial goal of this research has been a fully eddy-resolving, data assimilative global ocean prediction system with $1/16^\circ$ horizontal resolution and this is currently being transitioned to NAVO. An upgrade to the desired resolution of $1/32^\circ$ is planned for 2003.




Figure 4. The $1/16^\circ$ global SSH (sea surface height) forecast verification against the model with Topex/Poseidon and ERS-2 altimeter data assimilation for the period 1-31 January 1999. Shown is RMS error (cm) versus forecast length (days) for the Gulf Stream (left), the Kuroshio (middle) and the whole domain (right). The red curves are the Navy Layered Ocean Model forecasts; the blue curves are forecasts of persistence (i.e. no change from the initial state) and the black curve is from climatology.

Predictive Skill

Figure 3 depicts a sea surface height nowcast and 14 day forecast valid on 15 January 1999 from the 1/16° global system. An independent 1/8° SST analysis from the Modular Ocean Data Assimilation System (MODAS) is also shown for comparison. The SST analysis shows good correspondence between the Kuroshio pathway and some of the eddies seen in the simulated SSH (sea surface height) field. The global system consistently produces a skillful regional and basin-wide forecast of oceanic fronts and eddies. Figure 4 shows forecast verification from the period discussed above. Compared to forecasts of climatology and persistence (no change), the operational system shows forecast skill for more than one month over the entire model domain and Kuroshio region, and 21 day forecast skill in the Gulf Stream region in this instance.

The results have shown that altimeter data alone is sufficient to produce an accurate nowcast when a high resolution ocean model is in the loop to act as a dynamical interpolator to fill in the space-time gaps in the altimeter data. The simulation skill of the model also leads to skillful forecasts of a month or more in many regions.

The World Wide Web

To view real time Navy Layered Ocean Model results on the world wide web, start at the NAVOCEANO homepage (<http://www.navo.navy.mil>) and click on "operational products". In the sub-window click on "product search form". In the next sub-window click on the "product type view" button and in the lower window scroll to "Model Navy Layered Ocean Model" and highlight it. This causes several region indicators to appear in the upper window and these should be highlighted. Finally, click on the "Submit Query" button. The analysis and forecast plots and animations are then viewable. Near the top of the page is a hyperlink to aid in book marking references.

Acknowledgments

This work is a culmination of many projects over many years. Individuals who have made significant contributions toward this final operational product include Jay F. Shriver (NRL), A. Birol Kara (Sverdrup Technology Inc.), and Jean-Francois Cayula (Planning Systems Inc.). This is a contribution to the 6.2 Basin-scale Ocean Prediction System project funded by the Office of Naval Research under program element 62435N and to the 6.4 projects Large Scale Ocean Modeling and Ocean Data Assimilation funded by the Space and Naval Warfare Systems Command under program element 63207N. The numerical model was run on the Cray T3Es at the Naval Oceanographic Office, Stennis Space Center, Mississippi and at the U.S. Army Engineer Research and Development Center, Vicksburg, Mississippi. Both are part of the Defense Department's High Performance Computing Initiative. ■

References

- Hellerman, S. and M. Rosenstein, Normal Monthly Wind Stress Over the World Ocean with Error Estimates, *J. Phys. Oceanogr.*, 13, 1093-1104, 1983.
- Hurlburt, H.E. and P.J. Hogan, Impact of 1/8° to 1/64° Resolution on Gulf Stream Model-data Comparisons in Basin-scale Subtropical Atlantic Ocean Models, *Dyn. Atmos. Ocean.*, 32, 283-329, 2000.
- Wallcraft, A.J. and D.R. Moore, The NRL Layered Ocean Model, *Parallel Computing*, 23, 2227-2242, 1997.

**Check-out todays
aquatic remote
sensing news at:
WaterObserver.org**



Corporate Members

CORE

www.brook.edu

CSIRO

www.cossa.csiro.au

Defence Research Establishment Atlantic

www.drea.dnd.ca



EADS Deutschland GmbH

www.eads-nv.com



EARTHSAT

Earth Satellite Corporation

www.earthsat.com



Earth Search Sciences

www.earthsearch.com



ERDAS, Inc.

www.erdas.com

FFI

www.ffi.no

Finnish Institute of Marine Research

ice.fmi.fi/main_uk.html

Fugro GEOS

www.geos.com



GSE RENTALS LTD.

www.gserentals.co.uk

Hackensack Meadowlands Dev. Comm.

www.hmdc.state.nj.us



HOBILabs Inc.

www.hobilabs.com

Ocean Color Spectrum

Edited by: James Acker

NASA GDAAC, MD, USA, Raytheon ITSS
E-mail: acker@dacc.gsfc.nasa.gov

Contribution by: Daniel M. Palacios

Oregon State University, OR, USA

Karin A. Forney

Southwest Marine Fisheries Center, NMFS, NOAA, USA

Gene Feldman

NASA Goddard Space Flight Center, MD, USA

*This month's Ocean Color Spectrum features a host of interesting contributions from around the world. The featured contribution is provided by **Daniel M. Palacios**, with co-authors Karin A. Forney and Gene Feldman.*

Cetaceans (whales and dolphins) are at the top of the oceanic food chain. From this perspective, it is important to augment our understanding of the influence of oceanographic phenomena on cetacean patterns of occurrence. On the western side of the Galápagos Islands, strong environmental gradients result from the interaction between tropical and equatorial surface waters with the upwelled waters of the Equatorial Undercurrent. GalCet2K was a cruise designed to investigate mesoscale physical/biological interactions influencing the distribution and abundance of cetaceans in this region. The 15 day survey took place 5 to 19 April 2000 aboard R/V Odyssey, as a collaborative effort between the authors and the Ocean Alliance, a non-profit whale research and conservation organization (www.oceanalliance.org).

In conducting habitat assessments for cetaceans, a compromise was made between the requirements necessary to obtain precise estimates of abundance, and an environmental sampling design that will resolve the scales of interest. This is complicated by the fact that, at the higher trophic levels, species-environment relationships are a function of complex trophic and life-history dynamics that are not easily measured. Therefore, we must

rely on proxy variables that can be readily measured within the scope of a typical cruise. Our approach consisted of an intensive visual survey for cetaceans along the track of the vessel, while stopping two to three times per day to sample the upper 150

m of the water column with a small CTD instrument with an attached fluorometer. Along-track measurements of sonar intensity were also collected at two frequencies (50 and 250 kHz), as a bulk index of biomass of zooplankton and nekton (which are potential

Photo of dolphins. A school of short-beaked common dolphins, the most abundant small cetacean in offshore waters of the Galápagos Archipelago. The typical size of these schools is about 270 animals, but they can aggregate in groups of 3,000 or more.

Photo courtesy of Daniel Palacios




Figure 1. Water column zooplankton/nekton biomass index integrated to 150 m, as derived from sonar returns (50 plus 250 kHz). Scale is arbitrary. Dotted white line depicts measurement locations and red contour indicates a biomass index of one.